

# Absolute Extrema and the Extreme Value Theorem


(covers Stewart 4.1)

Calculus can help us find the biggest and smallest values of functions, if they exist.

Suppose we have a function  $f$  with domain  $D$ .

$f$  has an **absolute maximum** value at a point  $c \in D$  if  $f(c) \geq f(x)$  for all  $x \in D$ .


$f$  has an **absolute minimum** value at a point  $c \in D$  if  $f(c) \leq f(x)$  for all  $x \in D$ .



Absolute maxima and minima are also called global maxima and minima.

### Ex 1.


What are the absolute max and absolute min (if any) of  $y = x^2$  with the following domains?



Domain:  $(-\infty, \infty)$

Abs max: \_\_\_\_\_


Abs min: \_\_\_\_\_



Domain:  $[0, 2]$

Abs max: \_\_\_\_\_


Abs min: \_\_\_\_\_



Domain:  $(0, 2]$

Abs max: \_\_\_\_\_

Abs min: \_\_\_\_\_



Domain:  $(0, 2)$





Abs max: \_\_\_\_\_

Abs min: \_\_\_\_\_


Q: When are we guaranteed to have an absolute max and min?

A: The **Extreme Value Theorem** says  $f$  must be continuous on a closed interval:

If  $f(x)$  is continuous on a closed interval  $[a, b]$ , then it will have an absolute max and min on  $[a, b]$ .




To the right is an example of a function defined on the closed interval  $[0, 1]$ , with no absolute max. Because the function is discontinuous, the Extreme Value Theorem doesn't apply, so we're not guaranteed an absolute max.




**How to find absolute extrema on a closed interval**

1. Evaluate  $f$  at all critical numbers and endpoints.
2. Take the largest and smallest of these values.

**Ex 2.**

Find the absolute maximum and minimum values of  $f(x) = 10x(2 - \ln x)$  on the interval  $[1, e^2]$ .

**Ex 3.**

Find the absolute maximum and minimum values of  $f(x) = x^{2/3}$  on the interval  $[-2, 3]$ .

